	NAMBTHAFF
CTIES SUBCLASS	BA
G.F.G.	Q3VO7199A

Docket: 23623-7043; Ser. 097436,513; Filed 17709799

McCtreen, Doyle, Brown & Enersen, LLP – (415) 393-2000

Invertible: Chemically Modified Mutant Serine Hydrolases Show Improved Catalytic Activity and Chiral Selectivity

Fig. 1

APPROVED	7.6 F	7
ВУ	CLASS [.J: -, 45S
ORAFISMO:		Ĭ

McCutchen, Doyle, Brown & Enersen, LLP – (415) 393-2000
Invent John Bryan Jones, et al.

Title: Chemically Modified Mutant Serine Hydrolases Show Improved Catalytic Activity and Chiral Selectivity

Scheme 1. Modification of SBL mutants with Chiral Auxiliaries.

$$R = S \xrightarrow{QR^{1}} (R)-a R^{1} = Me$$

$$(R)-b R^{1} = H$$

$$R = S \xrightarrow{QR^{1}} C R^{1} = H$$

$$(R)-b R^{1} = H$$

$$R = S \xrightarrow{QR^{1}} C R^{1} = H$$

$$(R)-d R^{1} = iPr$$


$$(R)-e R^{1} = Ph$$

$$(R)-f R^{1} = Bn$$

$$R = S$$
 N
 O
 (R) - g
 $R^1 = Ph$
 $R = S$
 N
 O
 (R) - i

The corresponding (S) MTS ligands follow the same code scheme (i.e. (S)-a, (S)-b, (S)-d, (S)-e, (S)-f, (Ś)-g, (S)-h, (S)-i).

Scheme 2. Synthesis of Mandelate-based Ligands




Reagents: (i) Me₂SO₄, NaOH, H₂O, 37%; (ii) MeOH, H⁺; (iii) MOM-Cl, CH₂Cl₂, Et₃N (90% 2 steps); (iv) For (R)-3: BH₃, THF, 82%; For (R)-5: LiBH₄, THF, 97%; (v) MeSO₂Cl, CH₂Cl₂, Et₃N; For (R)-8: 100%; (vi) LiBr, acetone; For (R)-10: 84%; For (R)-11: 78% 2 steps; (vii) NaSSO₂CH₃, DMF; For (R)-12: 61%; (viii) TFA, H₂O, 82%.

Fig. 3

McCutchen, Doyle, Brown & Enersen, LLP – (415) 393-2000
Inventors: John Jones, et al.
Title: Chemically Modified Mutant Serine Hydrolases Show Improved Catalytic Activity and Chiral Selectivity

Scheme 3. Synthesis of Oxazolidinone-based Ligands



Reagents: (i) KOH, DMSO, Br (CH₂)_nBr; (ii) NaSSo₂CH₃, DMF.

McCutches Doyle, Brown & Enersen, LLP – (415) 393-2000 Inventors in Bryan Jones, et al.

Title: Chemically Modified Mutant Serine Hydrolases Show Improved Catalytic Activity and Chiral Selectivity

Scheme 4. Synthesis of Indanol-based Ligands

H₂N OH
(i)
(R)-23
(R)-24

(ii)
(R)-25 R = Br
(R)-11 R =
$$SSO_2CH_3$$

Reagents: (i) triphosgene, CH_2CI_2 , Et_3N , 100%; (ii) KOH, DMSO, $Br(CH_2)_3Br$; (iii) $NaSSO_2CH_3$, DMF.

Fig. 5

McCutchen Doyle, Brown & Enersen, LLP – (415) 393-2000 Invento Intellet Chemically Modified Mutant Serine Hydrolases Show Improved Catalytic Activity and Chiral Selectivity





Fig. 6A



APPRIORED D.G. F. 3.

McCutch Doyle, Brown & Enersen, LLP – (415) 393-2000 Invento Interest of Chemically Modified Mutant Serine Hydrolases Show Improved Catalytic Activity and Chiral Selectivity




Fig. 6C